

Samo Stanič, journal publications in 1998

References

- [1] P. Abreu *et al.* [DELPHI Collaboration], “A search for heavy stable and longlived squarks and sleptons in e^+e^- collisions at energies from 130 GeV to 183 GeV,” Phys. Lett. B **444**, 491 (1998) [arXiv:hep-ex/9811007].
- [2] P. Abreu *et al.* [DELPHI Collaboration], “Two-particle angular correlations in e^+e^- interactions compared with QCD predictions,” Phys. Lett. B **440**, 203 (1998).
- [3] P. Abreu *et al.* [DELPHI Collaboration], “A search for η_c ’ production in photon photon fusion at LEP,” Phys. Lett. B **441**, 479 (1998).
- [4] P. Abreu *et al.* [DELPHI Collaboration], “Measurement of $|V_{cs}|$ using W decays at LEP2,” Phys. Lett. B **439**, 209 (1998).
- [5] P. Abreu *et al.* [DELPHI Collaboration], “ π^\pm, K^\pm, p and \bar{p} production in $Z^0 \rightarrow q\bar{q}$, $Z^0 \rightarrow b\bar{b}$, $Z^0 \rightarrow u\bar{u}, d\bar{d}, s\bar{s}$,” Eur. Phys. J. C **5**, 585 (1998).
- [6] P. Abreu *et al.* [DELPHI Collaboration], “Measurement of the $e^+e^- \rightarrow \gamma\gamma(\gamma)$ cross section at the LEP energies,” Phys. Lett. B **433**, 429 (1998).
- [7] P. Abreu *et al.* [DELPHI Collaboration], “Measurement of the charged particle multiplicity of weakly decaying B hadrons,” Phys. Lett. B **425**, 399 (1998).
- [8] P. Abreu *et al.* [DELPHI Collaboration], “First evidence for a charm radial excitation, D^{*} ,” Phys. Lett. B **426**, 231 (1998).
- [9] P. Abreu *et al.* [DELPHI Collaboration], “Investigation of the splitting of quark and gluon jets,” Eur. Phys. J. C **4**, 1 (1998).
- [10] P. Abreu *et al.* [DELPHI Collaboration], “A Study of the hadronic resonance structure in the decay $\tau \rightarrow 3\pi + \nu_\tau$,” Phys. Lett. B **426**, 411 (1998).
- [11] P. Abreu *et al.* [DELPHI Collaboration], “Measurement of the inclusive charmless and double-charm B branching ratios,” Phys. Lett. B **426**, 193 (1998).
- [12] P. Abreu *et al.* [DELPHI Collaboration], “Measurement of trilinear gauge couplings in e^+e^- collisions at 161 GeV and 172 GeV,” Phys. Lett. B **423**, 194 (1998).
- [13] P. Abreu *et al.* [DELPHI Collaboration], “Measurement of the W pair cross-section and of the W mass in e^+e^- interactions at 172 GeV,” Eur. Phys. J. C **2**, 581 (1998).
- [14] P. Abreu *et al.* [DELPHI Collaboration], “Search for charged Higgs bosons in e^+e^- collisions at $\sqrt{s} = 172$ GeV,” Phys. Lett. B **420**, 140 (1998).
- [15] P. Abreu *et al.* [DELPHI Collaboration], “ m_b at M_Z ,” Phys. Lett. B **418**, 430 (1998).
- [16] P. Abreu *et al.* [DELPHI Collaboration], “Charged particle multiplicity in $e^+e^- \rightarrow q\bar{q}$ events at 161 GeV and 172 GeV and from the decay of the W boson,” Phys. Lett. B **416**, 233 (1998).
- [17] P. Abreu *et al.* [DELPHI Collaboration], “Search for charginos, neutralinos and gravitinos at LEP,” Eur. Phys. J. C **1**, 1 (1998).

- [18] P. Abreu *et al.* [DELPHI Collaboration], “Search for neutral and charged Higgs bosons in e^+e^- collisions at $\sqrt{s} = 161$ GeV and 172 GeV,” *Eur. Phys. J. C* **2**, 1 (1998).
- [19] P. Abreu *et al.* [DELPHI Collaboration], “Rapidity correlations in Lambda baryon and proton production in hadronic Z^0 decays,” *Phys. Lett. B* **416**, 247 (1998).