

References

- [1] S. Mori *et al.*, “The Belle detector,” Nucl. Instrum. Meth. A **479** 117 (2002).
- [2] K. Abe *et al.*, “Observation of color-suppressed $\bar{B}^0 \rightarrow D^0\pi^0$, $D^{*0}\pi^0$, $D^0\eta$ and $D^0\omega$ decays,” Phys. Rev. Lett. **88** 052002 (2002).
- [3] K. Abe *et al.*, “Observation of the decay $B \rightarrow K\mu^+\mu^-$,” Phys. Rev. Lett. **88** 021801 (2002).
- [4] K. Abe *et al.*, “Production of prompt charmonia in e^+e^- annihilation at $\sqrt{s} = 10.6$ GeV,” Phys. Rev. Lett. **88** 052001 (2002).
- [5] K. Abe *et al.*, “A measurement of lifetime difference in D^0 meson decays,” Phys. Rev. Lett. **88** 162001 (2002).
- [6] K. Abe *et al.*, “Observation of Cabibbo-suppressed and W exchange Λ_c^+ baryon decays,” Phys. Lett. B **524** 33 (2002).
- [7] K. Abe *et al.*, “Determination of $|V_{cb}|$ using the semileptonic decay $\bar{B}^0 \rightarrow D^{*+}e^-\bar{\nu}$,” Phys. Lett. B **526** 247 (2002).
- [8] K. Abe *et al.*, “Observation of $B^+ \rightarrow \chi_c^0 K^+$,” Phys. Rev. Lett. **88** 031802 (2002).
- [9] K. Abe *et al.*, “Measurement of $\text{Br}(\bar{B}^0 \rightarrow D^+l^-\bar{\nu})$ and determination of $|V_{cb}|$,” Phys. Lett. B **526** 258 (2002).
- [10] K. Abe *et al.*, “Study of three-body charmless B decays,” Phys. Rev. D **65** 092005 (2002).
- [11] K. Abe *et al.*, “Precise measurement of B meson lifetimes with hadronic decay final states,” Phys. Rev. Lett. **88** 171801 (2002).
- [12] K. Abe *et al.*, “Observation of $B^\pm \rightarrow p\bar{p}K^\pm$,” Phys. Rev. Lett. **88** 181803 (2002).
- [13] K. Abe *et al.*, “Observation of mixing-induced CP violation in the neutral B meson system,” Phys. Rev. D **66** 032007 (2002).
- [14] K. Abe *et al.*, “Observation of χ_{c2} production in B-meson decay,” Phys. Rev. Lett. **89** 011803 (2002).
- [15] K. Abe *et al.*, “Search for charmless two-body baryonic decays of B mesons,” Phys. Rev. D **65** 091103 (2002).
- [16] K. Abe *et al.*, “Measurement of CP-violating asymmetries in $B^0 \rightarrow \pi^+\pi^-$ decays,” Phys. Rev. Lett. **89**, 071801 (2002).
- [17] K. Abe *et al.*, “Measurements of branching fractions and decay amplitudes in $B \rightarrow J/\Psi K^*$ decays,” Phys. Lett. B **538** 11 (2002).
- [18] S. Nishida *et al.*, “Radiative B meson decays into $K\pi\gamma$ and $K\pi\pi\gamma$ final states,” Phys. Rev. Lett. **89**, 231801 (2002).
- [19] K. Abe *et al.*, “Observation of $\bar{B}^0 \rightarrow D^{(*)0}p\bar{p}$,” Phys. Rev. Lett. **89** 151802 (2002).
- [20] R. Abe *et al.*, “Status Of The Belle SVD Detector,” Nucl. Instrum. Meth. A **478**, 296 (2002).
- [21] K. Abe *et al.*, “Measurement of χ_{c2} production in two-photon collisions,” Phys. Lett. B **540** 33 (2002).

- [22] K. Abe *et al.*, “Observation of double $c\bar{c}$ production in e^+e^- annihilation at \sqrt{s} 10.6 GeV,” Phys. Rev. Lett. **89** 142001 (2002).
- [23] S. K. Choi *et al.*, “Observation of the $\eta_c(2s)$ in exclusive $B \rightarrow K K_s K^- \pi^+$ decays,” Phys. Rev. Lett. **89** 102001 (2002). [Erratum-ibid. **89** 129901 (2002)].
- [24] K. Abe *et al.*, “Observation of the decay $B^0 \rightarrow D^\pm D^{*\mp}$,” Phys. Rev. Lett. **89** 122001 (2002).
- [25] A. Gordon *et al.*, “Study of $B \rightarrow \rho \pi$ decays at Belle,” Phys. Lett. B **542**, 183 (2002).
- [26] R. S. Lu *et al.*, “Observation of $B^\pm \rightarrow \omega K^\pm$ decay,” Phys. Rev. Lett. **89**, 191801 (2002).
- [27] T. Tomura *et al.*, “Measurement of the oscillation frequency for $B^0 - \bar{B}^0$ mixing using hadronic B^0 decays,” Phys. Lett. B **542**, 207 (2002).
- [28] K. F. Chen *et al.*, “Measurement of CP-violating parameters in $B \rightarrow \eta' K$ decays,” Phys. Lett. B **546** 196 (2002).
- [29] A. Drutskoy *et al.*, “Observation of $B \rightarrow D^{(*)} K^- K^{(*)0}$ decays,” Phys. Lett. B **542**, 171 (2002).
- [30] P. Krokovny *et al.*, “Observation of $D_{s^+} K^-$ and evidence for $D_{s^+} \pi^-$ final states in neutral B decays,” Phys. Rev. Lett. **89**, 231804 (2002).
- [31] B. C. K. Casey *et al.*, “Charmless hadronic two-body B meson decays,” Phys. Rev. D **66**, 092002 (2002).
- [32] K. Abe *et al.*, “An improved measurement of mixing-induced CP violation in the neutral B meson system,” Phys. Rev. D **66** 071102 (2002).
- [33] K. Abe *et al.*, “Measurement of the inclusive semileptonic branching fraction of B mesons and $|V_{cb}|$,” Phys. Lett. B **547** 181 (2002).
- [34] N. Gabyshev *et al.*, “Study of exclusive B decays to charmed baryons at Belle,” Phys. Rev. D **66**, 091102 (2002).
- [35] K. Hara *et al.*, “Measurement Of The $B^0 \bar{B}^0$ Mixing Parameter Δm_d Using Semileptonic B^0 Decays,” Phys. Rev. Lett. **89**, 251803 (2002).