

References

- [1] L. Ruckman, G. Varner, S. Stanič, A. Koga and T. Tsuboyama, “Development and implementation of a readout module for radiation-sensing field-effect transistors,” IEEE Trans. Nucl. Sci. **53** (2006) 2452.
- [2] G. Varner *et al.*, “Development of the Continuous Acquisition Pixel (CAP) sensor for high luminosity lepton colliders,” Nucl. Instrum. Meth. A **565**, 126 (2006).
- [3] J. Dragic *et al.*, “Measurement of branching fraction and direct CP asymmetry in $B^0 \rightarrow \rho^0 \pi^0$ decays,” Phys. Rev. D **73**, 111105 (2006).
- [4] G. Gokhroo *et al.*, “Observation of a near-threshold $D^0 \bar{D}^0 \pi^0$ enhancement in $B \rightarrow D^0 \bar{D}^0 \pi^0 K$ decay,” Phys. Rev. Lett. **97** (2006) 162002.
- [5] R. Chistov *et al.*, “Observation of new states decaying into $\Lambda_c^+ K^- \pi^+$ and $\Lambda_c^+ K_S^0 \pi^-$,” Phys. Rev. Lett. **97** (2006) 162001.
- [6] C. H. Wu *et al.*, “Study of $J/\psi \rightarrow p\bar{p}$, $\Lambda\bar{\Lambda}$ and observation of $\eta_c \rightarrow \Lambda\bar{\Lambda}$ at Belle,” Phys. Rev. Lett. **97** (2006) 162003.
- [7] Y. Miyazaki *et al.*, “Search for lepton flavor violating τ^- decays with a K_S^0 meson,” Phys. Lett. B **639** (2006) 159.
- [8] F. Fang *et al.*, “Search for the h_c meson in $B^\pm \rightarrow h_c K^\pm$,” Phys. Rev. D **74**, 012007 (2006).
- [9] A. Poluektov *et al.*, “Measurement of ϕ_{33} with Dalitz plot analysis of $B^+ \rightarrow D^{(*)} K^{(*)}$ decay,” Phys. Rev. D **73**, 112009 (2006).
- [10] L. Widhalm *et al.*, “Measurement of $D^0 \rightarrow \pi l \nu (K l \nu)$ form factors and absolute branching fractions,” Phys. Rev. Lett. **97** (2006) 061804.
- [11] F. J. Ronga *et al.*, “Measurements of CP violation in $B^0 \rightarrow D^{*-} \pi^+$ and $B^0 \rightarrow D^- \pi^+$ decays,” Phys. Rev. D **73**, 092003 (2006).
- [12] Y. Yusa *et al.*, “Search for neutrinoless decays $\tau \rightarrow l h h$ and $\tau \rightarrow l V^0$,” Phys. Lett. B **640** (2006) 138.
- [13] A. Ishikawa *et al.*, “Measurement of forward-backward asymmetry and Wilson coefficients in $B \rightarrow K^* l^+ l^-$,” Phys. Rev. Lett. **96**, 251801 (2006).
- [14] J. Schumann *et al.*, “Evidence for $B \rightarrow \eta' \pi$ and improved measurements for $B \rightarrow \eta' K$,” Phys. Rev. Lett. **97** (2006) 061802.
- [15] A. Somov *et al.*, “Measurement of the branching fraction, polarization, and CP asymmetry for $B^0 \rightarrow \rho^+ \rho^-$ decays, and determination of the CKM phase ϕ_2 ,” Phys. Rev. Lett. **96**, 171801 (2006).

- [16] L. M. Zhang *et al.*, “Improved constraints on $D^0 - \bar{D}^0$ mixing in $D^0 \rightarrow K^+\pi^-$ decays at Belle,” Phys. Rev. Lett. **96**, 151801 (2006).
- [17] Z. Natkaniec *et al.*, “Status of the Belle silicon vertex detector,” Nucl. Instrum. Meth. A **560**, 1 (2006).
- [18] A. Garmash *et al.*, “Evidence for large direct CP violation in $B^\pm \rightarrow \rho(770)^0 K^\pm$ from analysis of the three-body charmless $B^\pm \rightarrow K^\pm \pi^\pm \pi^\mp$ decay,” Phys. Rev. Lett. **96**, 251803 (2006).
- [19] K. Abe *et al.*, “Measurement of azimuthal asymmetries in inclusive production of hadron pairs in e^+e^- annihilation at Belle,” Phys. Rev. Lett. **96**, 232002 (2006).
- [20] K. Abe *et al.*, “Observation of $b \rightarrow d\gamma$ and determination of $|V_{td}/V_{ts}|$,” Phys. Rev. Lett. **96**, 221601 (2006).
- [21] E. Nakano *et al.*, “Charge asymmetry of same-sign dileptons in $B^0 - \bar{B}^0$ mixing,” Phys. Rev. D **73**, 112002 (2006).
- [22] K. Abe *et al.*, “Study of $B^\pm \rightarrow D_{(CP)}K^\pm$ and $D_{(CP)}^*K^\pm$ decays,” Phys. Rev. D **73** (2006) 051106.
- [23] S. Uehara *et al.*, “Observation of a χ'_{c2} candidate in $\gamma\gamma \rightarrow D\bar{D}$ production at Belle,” Phys. Rev. Lett. **96** (2006) 082003.
- [24] Y. Miyazaki *et al.* [Belle Collaboration], “Search for Lepton Flavor Violation in τ decays at B-factories,” PoS **HEP2005** (2006) 317.
- [25] Y. Miyazaki *et al.*, “Search for lepton and baryon number violating τ^- decays into $\bar{\Lambda}\pi^-$ and $\Lambda\pi^-$,” Phys. Lett. B **632** (2006) 51.
- [26] N. Soni *et al.*, “Measurement of branching fractions for $B \rightarrow \chi_{c1(2)}K(K^*)$ at Belle,” Phys. Lett. B **634** (2006) 155.
- [27] K. Abe *et al.*, “Search for the decay $B^0 \rightarrow \gamma\gamma$,” Phys. Rev. D **73** (2006) 051107.
- [28] K. Abe *et al.*, “Search for the $\Theta(1540)^+$ pentaquark using kaon secondary interactions at Belle,” Phys. Lett. B **632** (2006) 173.
- [29] R. Seuster *et al.*, “Charm hadrons from fragmentation and B decays in e^+e^- annihilation at $\sqrt{s} = 10.6\text{GeV}$,” Phys. Rev. D **73** (2006) 032002.
- [30] S. Blyth *et al.*, “Improved measurements of color-suppressed decays $\bar{B}^0 \rightarrow D^0\pi^0, D^0\eta, D^0\omega, D^{*0}\pi^0, D^{*0}\eta$ and $D^{*0}\omega$,” Phys. Rev. D **74** (2006) 092002.
- [31] K. Abe *et al.*, “The first observation of $\tau^\pm \rightarrow \Phi K^\pm \nu$ decay,” Phys. Lett. B **643** (2006) 5.